



Bright Futures Parent Handout

2 Year Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

Your Talking Child

- Talk about and describe pictures in books and the things you see and hear together.
- Parent-child play, where the child leads, is the best way to help toddlers learn to talk.
- Read to your child every day.
- Your child may love hearing the same story over and over.
- Ask your child to point to things as you read.
- Stop a story to let your child make an animal sound or finish a part of the story.
- Use correct language; be a good model for your child.
- Talk slowly and remember that it may take a while for your child to respond.

ASSESSMENT OF LANGUAGE DEVELOPMENT

Your Child and TV

- It is better for toddlers to play than watch TV.
- Limit TV to 1–2 hours or less each day.
- Watch TV together and discuss what you see and think.
- Be careful about the programs and advertising your young child sees.
- Do other activities with your child such as reading, playing games, and singing.
- Be active together as a family. Make sure your child is active at home, at child care, and with sitters.

TELEVISION VIEWING

Safety

- Be sure your child's car safety seat is correctly installed in the back seat of all vehicles.
- All children 2 years or older should use a forward-facing car safety seat with a harness for as long as possible, up to the highest weight or height allowed by their car safety seat's manufacturer.

SAFETY

SAFETY

- Everyone should wear a seat belt in the car. Do not start the vehicle until everyone is buckled up.
- Never leave your child alone in your home or yard, especially near cars, without a mature adult in charge.
- When backing out of the garage or driving in the driveway, have another adult hold your child a safe distance away so he is not run over.
- Keep your child away from moving machines, lawn mowers, streets, moving garage doors, and driveways.
- Have your child wear a good-fitting helmet on bikes and trikes.
- If you have a gun, store it unloaded and locked with the ammunition locked separately from the gun.

Toilet Training

- Signs of being ready for toilet training
 - Dry for 2 hours
 - Knows if she is wet or dry
 - Can pull pants down and up
 - Wants to learn
 - Can tell you if she is going to have a bowel movement
- Plan for toilet breaks often. Children use the toilet as many as 10 times each day.
- Help your child wash her hands after toileting and diaper changes and before meals.
- Clean potty chairs after every use.
- Teach your child to cough or sneeze into her shoulder. Use a tissue to wipe her nose.
- Take the child to choose underwear when she feels ready to do so.

TOILET TRAINING

How Your Child Behaves

- Praise your child for behaving well.
- It is normal for your child to protest being away from you or meeting new people.
- Listen to your child and treat him with respect. Expect others to do as well.
- Play with your child each day, joining in things the child likes to do.
- Hug and hold your child often.
- Give your child choices between 2 good things in snacks, books, or toys.
- Help your child express his feelings and name them.
- Help your child play with other children, but do not expect sharing.
- Never make fun of the child's fears or allow others to scare your child.
- Watch how your child responds to new people or situations.

TEMPERAMENT AND BEHAVIOR

What to Expect at Your Child's 3 Year Visit We will talk about

- Your talking child
- Getting ready for preschool
- Family activities
- Home and car safety
- Getting along with other children

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics, Updated 8/11. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.

Childhood Immunization Schedule

Birth

Hepatitis B

2 Week

No regularly scheduled vaccines

2 Month

DTaP

HIB

Polio (IPV)

Pneumococcal Conjugate (PCV-13)

Hepatitis B

Rotavirus

4 Month

DTaP

HIB

Polio (IPV)

Pneumococcal Conjugate (PCV-13)

Rotavirus

6 Month

DTaP

HIB

Polio (IPV)

Pneumococcal Conjugate (PCV-13)

Rotavirus

9 Month

Hepatitis B

12 Month

Hepatitis A

Pneumococcal Conjugate (PCV-13)

15 Month

DTaP

HIB

Polio (IPV)

MMR

Varicella

18 Month

No regularly scheduled vaccines

2 Year

Hepatitis A

30 Month

No regularly scheduled vaccines

3 & 4 Year

No regularly scheduled vaccines

5 Year

DTaP

Polio (IPV)

MMR

Varicella

7 & 8 Year

No regularly scheduled vaccines

9 & 10 Year

HPV

11 & 12 Year

TDaP

Meningococcal (ACWY)

HPV

13, 14, & 15 Year

No regularly scheduled vaccines

16 & 17 Year

Meningococcal (ACWY)

Meningococcal B

Hepatitis A Vaccine:

What You Need to Know

Many vaccine information statements are available in Spanish and other languages. See www.immunize.org/vis

Hojas de información sobre vacunas están disponibles en español y en muchos otros idiomas. Visite www.immunize.org/vis

1. Why get vaccinated?

Hepatitis A vaccine can prevent **hepatitis A**.

Hepatitis A is a serious liver disease. It is usually spread through close, personal contact with an infected person or when a person unknowingly ingests the virus from objects, food, or drinks that are contaminated by small amounts of stool (poop) from an infected person.

Most adults with hepatitis A have symptoms, including fatigue, low appetite, stomach pain, nausea, and jaundice (yellow skin or eyes, dark urine, light-colored bowel movements). Most children less than 6 years of age do not have symptoms.

A person infected with hepatitis A can transmit the disease to other people even if he or she does not have any symptoms of the disease.

Most people who get hepatitis A feel sick for several weeks, but they usually recover completely and do not have lasting liver damage. In rare cases, hepatitis A can cause liver failure and death; this is more common in people older than 50 years and in people with other liver diseases.

Hepatitis A vaccine has made this disease much less common in the United States. However, outbreaks of hepatitis A among unvaccinated people still happen.

2. Hepatitis A vaccine

Children need 2 doses of hepatitis A vaccine:

- First dose: 12 through 23 months of age
- Second dose: at least 6 months after the first dose

Infants 6 through 11 months old traveling outside the United States when protection against hepatitis A is recommended should receive 1 dose of hepatitis A vaccine. These children should still get 2 additional doses at the recommended ages for long-lasting protection.

Older children and adolescents 2 through 18 years of age who were not vaccinated previously should be vaccinated.

Adults who were not vaccinated previously and want to be protected against hepatitis A can also get the vaccine.

Hepatitis A vaccine is also recommended for the following people:

- International travelers
- Men who have sexual contact with other men
- People who use injection or non-injection drugs
- People who have occupational risk for infection
- People who anticipate close contact with an international adoptee
- People experiencing homelessness
- People with HIV
- People with chronic liver disease

In addition, a person who has not previously received hepatitis A vaccine and who has direct contact with someone with hepatitis A should get hepatitis A vaccine as soon as possible and within 2 weeks after exposure.

Hepatitis A vaccine may be given at the same time as other vaccines.



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

3. Talk with your health care provider

Tell your vaccination provider if the person getting the vaccine:

- Has had an **allergic reaction after a previous dose of hepatitis A vaccine**, or has any **severe, life-threatening allergies**

In some cases, your health care provider may decide to postpone hepatitis A vaccination until a future visit.

Pregnant or breastfeeding people should be vaccinated if they are at risk for getting hepatitis A. Pregnancy or breastfeeding are not reasons to avoid hepatitis A vaccination.

People with minor illnesses, such as a cold, may be vaccinated. People who are moderately or severely ill should usually wait until they recover before getting hepatitis A vaccine.

Your health care provider can give you more information.

4. Risks of a vaccine reaction

- Soreness or redness where the shot is given, fever, headache, tiredness, or loss of appetite can happen after hepatitis A vaccination.

People sometimes faint after medical procedures, including vaccination. Tell your provider if you feel dizzy or have vision changes or ringing in the ears.

As with any medicine, there is a very remote chance of a vaccine causing a severe allergic reaction, other serious injury, or death.

5. What if there is a serious problem?

An allergic reaction could occur after the vaccinated person leaves the clinic. If you see signs of a severe allergic reaction (hives, swelling of the face and throat, difficulty breathing, a fast heartbeat, dizziness, or weakness), call **9-1-1** and get the person to the nearest hospital.

For other signs that concern you, call your health care provider.

Adverse reactions should be reported to the Vaccine Adverse Event Reporting System (VAERS). Your health care provider will usually file this report, or you can do it yourself. Visit the VAERS website at www.vaers.hhs.gov or call **1-800-822-7967**. *VAERS is only for reporting reactions, and VAERS staff members do not give medical advice.*

6. The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) is a federal program that was created to compensate people who may have been injured by certain vaccines. Claims regarding alleged injury or death due to vaccination have a time limit for filing, which may be as short as two years. Visit the VICP website at www.hrsa.gov/vaccinecompensation or call **1-800-338-2382** to learn about the program and about filing a claim.

7. How can I learn more?

- Ask your health care provider.
- Call your local or state health department.
- Visit the website of the Food and Drug Administration (FDA) for vaccine package inserts and additional information at www.fda.gov/vaccines-blood-biologics/vaccines.
- Contact the Centers for Disease Control and Prevention (CDC):
 - Call **1-800-232-4636 (1-800-CDC-INFO)** or
 - Visit CDC's website at www.cdc.gov/vaccines.

